

## Cobalt-Catalyzed Regioselective Carboamidation of Alkynes with Imides Enabled by Cleavage of C–N and C–C Bonds

Xiang-Ting Min, Ding-Wei Ji, Hao Zheng, Bing-Zhi Chen, Yan-Cheng Hu, Boshun Wan, and Qing-An Chen\*



Cite This: *Org. Lett.* 2020, 22, 3386–3391



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

**ABSTRACT:** Through the oxidative addition of cobalt into the N–C(O) bond of phthalimide and the subsequent decarbonylation, we describe an efficient cobalt-catalyzed intermolecular decarbonylative carboamidation of alkynes. High regioselectivities have been achieved for unsymmetrical alkynes (including aryl–alkyl or aryl–aryl) to deliver polysubstituted isoquinolones. To facilitate step economy, a three-component decarbonylative carboamidation of alkynes with phthalic anhydrides and amines has been demonstrated using the current cobalt catalysis.




The activation of  $\sigma$ -bonds via oxidative addition with transition metals is a fundamental step in organometallic chemistry. Therefore, the development of novel transition metal catalytic systems for facilitating such a process is one of the leading edge efforts in catalysis.<sup>1</sup> Compared with the widely used noble metals, the earth-abundant first-row transition metals have gained particular attention in recent years owing to their relatively lower cost and toxicity<sup>2a,b,c</sup> as well as greater sustainability. Among them, cobalt catalysis has emerged as a powerful tool and substantial progress has been achieved in this area.<sup>2d,e</sup> For example, cobalt-catalyzed C(sp<sup>2</sup>)–H activation via oxidative addition has been well established (**Scheme 1a**, eq 1).<sup>3,4</sup> However, the oxidative addition of cobalt with C–C bonds is generally limited to strained rings, such as benzocyclobutenones<sup>5a</sup> and methylenecyclopropanes,<sup>5b,c</sup> and others<sup>6</sup> (**Scheme 1a**, eq 2). Furthermore, the C–N activation of amide under cobalt catalysis is rare. In 2017, Danoun and co-workers described the first cobalt-catalyzed activation of N-Boc-amides (twisted amides) to form esters for the first time<sup>7</sup> (**Scheme 1a**, eq 3). Despite advances in the transition metal catalyzed decarbonylation of amide,<sup>8</sup> the cobalt-catalyzed decarbonylative coupling of imides requiring both C–N and C–C bond cleavages remains undiscovered (**Scheme 1a**, eq 4).


On the other hand, catalytic multicomponent reaction, due to its rapid construction of multiple new bonds in one step, has received much attention over the past decades.<sup>9</sup> We envisioned phthalimide derivatives could be formed *in situ* from the condensation of phthalic anhydrides with amines (**Scheme 1b**). Subsequently, the aryl-cobalt intermediate generated from activating its C–N and C–C bonds will undergo migratory insertion into alkynes to give substituted isoquinolones (**Scheme 1b**). The challenges for this strategy include the following: (a) facile CO insertion into the C–Co bond makes it difficult to dissociate CO from cobalt;<sup>10,11</sup> (b) competitive

### Scheme 1. Activation of $\sigma$ -Bond via Cobalt Catalysis

a) Activation of  $\sigma$ -bond via oxidative addition with cobalt catalysis



b) This work: Cobalt-catalyzed decarbonylation of imides



decarbonylative homocoupling<sup>12</sup> and protolysis<sup>13</sup> due to production of water in the three-component reaction; (c) regiocontrol in the carboamidation of unsymmetrical alkynes.

Received: March 9, 2020

Published: April 10, 2020



Herein, we report a cobalt-catalyzed intermolecular decarbonylative carboamidation of alkynes to form substituted isoquinolones (**Scheme 1b**).

The two-component decarbonylative carboamidation of alkyne **2a** with phthalimide derivative **1a** was initially investigated (**Table 1**). After careful evaluation of the reaction

**Table 1. Selected Optimization Studies<sup>a</sup>**

entry	deviation from standard condition	3aa/3aa'	yield (3aa) <sup>b</sup>
1	<i>none</i>	14.3	92% <sup>c</sup>
2	without $\text{CoI}_2$	—	NR
3	without $\text{Ag}_2\text{CO}_3$	—	NR
4	without $\text{P}(\text{2-MeOC}_6\text{H}_4)_3$	4.0	28%
5	$\text{Co}(\text{PMe}_3)_4$ instead of $\text{CoI}_2$	—	NR
6	$\text{Co}_2(\text{CO})_8$ instead of $\text{CoI}_2$	—	NR
7	$\text{Co}(\text{PPh}_3)_3\text{Cl}$ instead of $\text{CoI}_2$	8.9	61%
8	$\text{Co}(\text{dppe})\text{Cl}_2$ instead of $\text{CoI}_2$	5.7	56%
9	$\text{Cp}^*\text{Co}(\text{CO})\text{I}_2$ instead of $\text{CoI}_2$	—	8%
10	$\text{PPh}_3$ instead of $\text{P}(\text{2-MeOC}_6\text{H}_4)_3$	5.6	35%
11	$\text{P}(\text{2-furyl})_3$ instead of $\text{P}(\text{2-MeOC}_6\text{H}_4)_3$	3.7	33%
12	Xantphos instead of $\text{P}(\text{2-MeOC}_6\text{H}_4)_3$	5.0	80%
13	$\text{Ag}_2\text{O}$ instead of $\text{Ag}_2\text{CO}_3$	8.3	86% <sup>c</sup>
14	$\text{AgOAc}$ instead of $\text{Ag}_2\text{CO}_3$	—	14%
15	TBHP instead of $\text{Ag}_2\text{CO}_3$	—	NR
16	$\text{AgOAc}$ and TBHP instead of $\text{Ag}_2\text{CO}_3$	8.0	52%


<sup>a</sup>All reactions were performed on 0.10 mmol scale. See Supporting Information for full details. <sup>b</sup>Determined by <sup>1</sup>H NMR analysis.

<sup>c</sup>Isolated yield.

conditions, isoquinolone **3aa** was formed in 92% yield with 14:1 (3aa/3aa') regioselectivity using  $\text{CoI}_2/\text{P}(\text{2-MeOC}_6\text{H}_4)_3$  as the metal/ligand combination and  $\text{Ag}_2\text{CO}_3$  as the oxidant (entry 1). Control experiments were subsequently conducted to understand the role of each reagent. In the absence of the cobalt catalyst or  $\text{Ag}_2\text{CO}_3$ , no desired product was observed (entries 2–3). Both the reactivity and regioselectivity decreased in the absence of  $\text{P}(\text{2-MeOC}_6\text{H}_4)_3$  (entry 4). Interestingly, the reaction did not proceed with  $\text{Co}^0$  precursors (entries 5–6). The use of a  $\text{Co}^{\text{I}}$  or  $\text{Co}^{\text{II}}$  precursor, such as  $\text{Co}(\text{PPh}_3)_3\text{Cl}$  and  $\text{Co}(\text{dppe})\text{Cl}_2$ , slightly lowered the yields (entries 7–8) (see Table S1). Probably due to the lack of available coordination sites, the employment of high valent  $\text{Cp}^*\text{Co}(\text{CO})\text{I}_2$  as the catalyst precursor gave only an 8% yield of **3aa** (entry 9). Other monodentate phosphine ligands, such as  $\text{PPh}_3$  or  $\text{P}(\text{2-furyl})_3$ , led to inferior results (entries 10–11). The bidentate ligand Xantphos could also promote this transformation but in slightly diminished yield (entry 12). The use of  $\text{Ag}_2\text{O}$  instead of  $\text{Ag}_2\text{CO}_3$  as the oxidant also gave isoquinolone **3aa** in a good yield (entry 13). However, the use of either  $\text{AgOAc}$  or TBHP as the oxidants significantly inhibited the reaction, thus suggesting both the oxidant and silver ion are important for generating the active intermediate (entries 14–16).

With the optimal conditions in hand, the substrate scope for the Co-catalyzed carboamidation of alkynes with phthalimides was investigated (**Scheme 2**). A variety of unsymmetrical aryl–alkyl alkynes underwent carboamidation smoothly to deliver


**Scheme 2. Scope of the Alkynes<sup>a</sup>**



<sup>a</sup>Isolated yield of the major regioselective isomer.


isoquinolone derivatives (**3aa**–**3af**) with high yields and regioselectivities. Cyclopropyl, ester, and alkoxy groups were well tolerated. For aryl–alkyl alkynes bearing indoles, good yields and excellent regioselectivities were observed (**3ag**–**3ah**). The regioselectivity for this unsymmetrical alkyne carboamidation was confirmed by X-ray analysis of **3ah**. A series of symmetrical diarylacetylenes were also compatible with this protocol, giving the corresponding products in good to excellent yields (**3ai**–**3an**). A heterocycle, such as thiophene, was well tolerated under the cobalt catalysis (**3ao**). It is interesting to note that, in the case of the unsymmetrical aryl–aryl alkyne with a slight difference in the electronic property of the two aryl substituents, the annulation delivered 3,4-diaryl substituted isoquinolone **3ap** with excellent regioselectivity (>20:1).

Next, selective decarbonylative annulation of phthalimide derivatives with alkynes that also bear phthalimide groups, which are widely embedded in pharmaceuticals and biologically active molecules such as thalidomide<sup>14</sup> were tested (**Scheme 3**). Selective C–N activation in the presence of four amide bonds on two structurally similar substrates (**1a** and **2q**–**2s**) may pose a challenge for the cobalt catalysis. Satisfactorily, the tested alkynes were compatible with this protocol, providing isoquinolones bearing phthalimide groups in high yields and regioselectivities (>20:1, **3aq**–**3ar**). Furthermore, isoquinolone **3as** with multiple N-heterocycles was isolated in moderate yield and regioselectivity. To demonstrate the scalability of this methodology, the reaction

**Scheme 3.** Selective Activation Among Four Amide Bonds<sup>a</sup>

of **1a** with **2q** was successfully performed on 2.0 mmol of starting materials, which afforded 0.862 g of **3aq** (85% yield).


After exploring the scope of the alkynes, an array of phthalimide derivatives were examined (Scheme 4). 3-Amino substituted phthalimides are prevalent motifs in drugs<sup>15</sup> and fluorescent probes;<sup>16a–c</sup> thus, it would be of interest to modify

**Scheme 4.** Scope of the Phthalimides<sup>a</sup>




the substrates of such motifs. The regioselective annulations of 3-amino substituted phthalimides **1b** and **1c** were successfully demonstrated. Moreover, it was found that the free NH on **1b** and **1c** plays an important role in the regiocontrol, because formation of its tautomer may hinder the oxidative addition of the adjacent carbonyl moiety with cobalt.<sup>16a,b,d</sup> The reaction of 3-chlorophthalimide **1d** with alkyne **2i** afforded cycloadduct **5di** in 60% yield, the product of which is difficult to synthesize via C–H activation owing to steric hindrance.<sup>17</sup> Phthalimides such as **1e**, **1f**, **1g**, **1h** were also compatible with the process, providing the corresponding products **4ei**, **4fi**, **4gi**, **4ga**, **4ha** in acceptable yields. 8-Amino-5-methoxyquinoline (MQ) can be used instead of 8-aminoquinoline for its ease of cleavage.<sup>18</sup>

Mechanistically, we sought to probe the aryl–cobalt intermediates resulting from the oxidative addition of a low-valent cobalt into the N–C(O) bond of imides and the subsequent decarbonylation (eqs 5–8). Interestingly, the

#### Decarbonylative homocoupling:




#### Protodecarbonylation:




decarbonylative homocoupling product **6a** can be obtained in 35% yield in the absence of alkynes.<sup>12</sup> The addition of stoichiometric H<sub>2</sub>O prevents the formation of **6a** (eq 5). When **1i** and **2a** were treated with the standard conditions, the corresponding cyclization product was not detected, and instead, decarbonylative product **6b** was obtained in 45% yield. The use of external water as an additive gave product **6b** in 90% yield (eq 6). Full incorporation of deuterium at the site of decarbonylation (**6c**) in the presence of D<sub>2</sub>O suggested that water was the proton source for this reaction (eq 7).<sup>13</sup> To gain insight into the role of the oxidant, the gas phase of the reaction mixture was analyzed by GC-MS (eq 8). Silver oxide (Ag<sub>2</sub>O) was used instead of Ag<sub>2</sub>CO<sub>3</sub> to avoid the background decarboxylation of Ag<sub>2</sub>CO<sub>3</sub> (eq 8). Carbon dioxide (CO<sub>2</sub>) was detected as the major gas component, which suggested that the oxidant played a role as a CO scavenger in the cobalt-catalyzed decarbonylation.

A proposed catalytic cycle is depicted in Scheme 5. First, Co(II) is disproportionated to active Co(I) species A.<sup>19</sup> Through an oxidative addition process, Co(I) species A activates the C–N bond of imides to form Co(III) intermediate B (from the perspective of ring strained, N–C activation precedes C–C activation; see the Supporting Information for details). A subsequent decarbonylation leads

**Scheme 5. Proposed Mechanism**

to the cobalt complex C. As a CO scavenger, the addition of an external oxidant facilitates the formation of active intermediate aryl-Co D. In the presence of alkyne 2, intermediate D undergoes coordination and migratory insertion of the alkyne to afford seven-membered cobaltacycle E. A final reductive elimination produces the isoquinolone products and regenerates Co(I) catalyst A. In the absence of alkyne 2, aryl-cobalt C undergoes a homocoupling pathway to yield compound 6a<sup>12</sup> or protolysis to yield compound 6c.<sup>13</sup>

A three-component decarbonylative carboamidation was accomplished by the use of bulk chemicals phthalic anhydride 7, 8-aminoquinoline 8, and alkynes 2 (Scheme 6). Overall, this three-component protocol cleaves four existing bonds while creating three additional new bonds in one pot. Gratifyingly, under the standard reaction conditions, a series of representative isoquinolones were obtained in moderate to high yields (3aa, 3ac, 3af, 3ai, 3ap, 3ah, 3aq, 4ha). Compared with the traditional C–H activation strategy, this three-

**Scheme 6. Three-Component Decarbonylative Carboamidation of Alkynes<sup>a</sup>**

<sup>a</sup>Isolated yield of major regioselective isomer.

component reaction provides a more convenient alternative to access such cyclic products.<sup>17</sup>

In summary, we have described an efficient Co-catalyzed intermolecular decarbonylative carboamidation of alkynes via the cleavage of C–N and C–C bonds of imides. The migratory insertion of aryl–cobalt species with unsymmetrical alkynes (including aryl–alkyl or aryl–aryl) occurs in high regiocontrol to deliver polysubstituted isoquinolones. A three-component Co-catalyzed decarbonylative carboamidation of alkynes with phthalic anhydrides and amines has been also demonstrated to increase step economy. Further application of this strategy to other amide decarbonylations and detailed mechanistic investigations are currently ongoing in our laboratory.

**■ ASSOCIATED CONTENT****SI Supporting Information**

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c00875>.

Experimental materials and procedures, NMR of compounds, and X-ray crystallographic analysis for compounds 3ap, 3ah, 3aq, and 4ba (PDF)

**Accession Codes**

CCDC 1974651, 1974659, 1974661, 1974675, and 1977034 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

**■ AUTHOR INFORMATION****Corresponding Author**

Qing-An Chen – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China; [orcid.org/0000-0002-9129-2656](https://orcid.org/0000-0002-9129-2656); Email: [qachen@dicp.ac.cn](mailto:qachen@dicp.ac.cn)

**Authors**

Xiang-Ting Min – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China; University of Chinese Academy of Sciences, Beijing 100049, China

Ding-Wei Ji – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China; University of Chinese Academy of Sciences, Beijing 100049, China

Hao Zheng – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China; University of Chinese Academy of Sciences, Beijing 100049, China

Bing-Zhi Chen – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China; University of Chinese Academy of Sciences, Beijing 100049, China

Yan-Cheng Hu – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China; [orcid.org/0000-0002-8731-4445](https://orcid.org/0000-0002-8731-4445)

Boshun Wan – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China; [orcid.org/0000-0002-7001-6214](https://orcid.org/0000-0002-7001-6214)

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.orglett.0c00875>

**Notes**

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We thank Prof. Yong-Gui Zhou (DICP), Prof. Martin Oestreich (TU Berlin), and Prof. Kevin G. M. Kou (UCR) for helpful discussions and manuscript revisions. Financial support from the National Natural Science Foundation of China (21971234, 21772194, 21572225), Liaoning Revitalization Talents Program (XLYC1807181), and the “Thousand Youth Talents Plan” is acknowledged.

## REFERENCES

- (1) (a) Chu, T.; Nikonov, G. I. Oxidative Addition and Reductive Elimination at Main-Group Element Centers. *Chem. Rev.* **2018**, *118*, 3608. (b) Kancherla, R.; Muralirajan, K.; Maity, B.; Zhu, C.; Krach, P. E.; Cavallo, L.; Rueping, M. Oxidative Addition to Palladium(0) Made Easy through Photoexcited-State Metal Catalysis: Experiment and Computation. *Angew. Chem., Int. Ed.* **2019**, *58*, 3412. (c) Uehling, M. R.; King, R. P.; Kraska, S. W.; Cernak, T.; Buchwald, S. L. Pharmaceutical Diversification via Palladium Oxidative Addition Complexes. *Science* **2019**, *363*, 405. (d) Xia, Z.; Corce, V.; Zhao, F.; Przybylski, C.; Espagne, A.; Jullien, L.; Le Saux, T.; Gimbert, Y.; Dossmann, H.; Mouries-Mansuy, V.; Ollivier, C.; Fensterbank, L. Photosensitized Oxidative Addition to Gold(I) Enables Alkynylative Cyclization of O-Alkylnylphenols with Iodoalkynes. *Nat. Chem.* **2019**, *11*, 797.
- (2) (a) Egorova, K. S.; Ananikov, V. P. Which Metals are Green for Catalysis? Comparison of the Toxicities of Ni, Cu, Fe, Pd, Pt, Rh, and Au Salts. *Angew. Chem., Int. Ed.* **2016**, *55*, 12150. (b) Li, J.; Tan, E.; Keller, N.; Chen, Y. H.; Zehetmaier, P. M.; Jakowitz, A. C.; Bein, T.; Knochel, P. Cobalt-Catalyzed Electrophilic Aminations with Anthranils: An Expedient Route to Condensed Quinolines. *J. Am. Chem. Soc.* **2019**, *141*, 98. (c) [LD<sub>50</sub> (CoCl<sub>2</sub>, rat oral) = 418 mg/kg; LD<sub>50</sub> (NiCl<sub>2</sub>, rat oral) = 186 mg/kg; LD<sub>50</sub> (PdCl<sub>2</sub>, rat oral) = 200 mg/kg]. (d) Pabst, T. P.; Obligacion, J. V.; Rochette, E.; Pappas, I.; Chirik, P. J. Cobalt-Catalyzed Borylation of Fluorinated Arenes: Thermodynamic Control of C(sp<sup>2</sup>)-H Oxidative Addition Results in Ortho-to-Fluorine Selectivity. *J. Am. Chem. Soc.* **2019**, *141*, 15378. (e) Wang, D.; Leng, X.; Ye, S.; Deng, L. Substrate Redox Non-Innocence Inducing Stepwise Oxidative Addition Reaction: Nitrosoarene C-N Bond Cleavage on Low-Coordinate Cobalt(0) Species. *J. Am. Chem. Soc.* **2019**, *141*, 7731.
- (3) (a) Gao, K.; Lee, P. S.; Fujita, T.; Yoshikai, N. Cobalt-Catalyzed Hydroarylation of Alkynes through Chelation-Assisted C-H Bond Activation. *J. Am. Chem. Soc.* **2010**, *132*, 12249. (b) Lee, P. S.; Fujita, T.; Yoshikai, N. Cobalt-Catalyzed, Room-Temperature Addition of Aromatic Imines to Alkynes via Directed C-H Bond Activation. *J. Am. Chem. Soc.* **2011**, *133*, 17283. (c) Ding, Z.; Yoshikai, N. Mild and Efficient C2-Alkenylation of Indoles with Alkynes Catalyzed by a Cobalt Complex. *Angew. Chem., Int. Ed.* **2012**, *51*, 4698. (d) Xu, W.; Yoshikai, N. N-H Imine as a Powerful Directing Group for Cobalt-Catalyzed Olefin Hydroarylation. *Angew. Chem., Int. Ed.* **2016**, *55*, 12731. (e) Sun, Q.; Yoshikai, N. Cobalt-Catalyzed Tandem Radical Cyclization/C-C Coupling Initiated by Directed C-H Activation. *Org. Lett.* **2019**, *21*, 5238.
- (4) (a) Fallon, B. J.; Garsi, J.-B.; Derat, E.; Amatore, M.; Aubert, C.; Petit, M. Synthesis of 1,2-Dihydropyridines Catalyzed by Well-Defined Low-Valent Cobalt Complexes: C-H Activation Made Simple. *ACS Catal.* **2015**, *5*, 7493. (b) Fallon, B. J.; Derat, E.; Amatore, M.; Aubert, C.; Chemla, F.; Ferreira, F.; Perez-Luna, A.; Petit, M. C-H Activation/Functionalization Catalyzed by Simple, Well-Defined Low-Valent Cobalt Complexes. *J. Am. Chem. Soc.* **2015**, *137*, 2448. (c) Fallon, B. J.; Derat, E.; Amatore, M.; Aubert, C.; Chemla, F.; Ferreira, F.; Perez-Luna, A.; Petit, M. C2-Alkylation and Alkenylation of Indoles Catalyzed by a Low-Valent Cobalt Complex in the Absence of Reductant. *Org. Lett.* **2016**, *18*, 2292.
- (5) For selected examples of benzocyclobutenones, see: (a) Zhu, Z.; Li, X.; Chen, S.; Chen, P. H.; Billett, B. A.; Huang, Z.; Dong, G. Cobalt-Catalyzed Intramolecular Alkyne/Benzo-cyclobutene Coupling: C-C Bond Cleavage via a Tetrahedral Dicobalt Intermediate. ACS Catal. **2018**, *8*, 845. (b) Kurahashi, T.; de Meijere, A. C-C Bond Activation by Octacarbonyldicobalt: [3 + 1] Cycloadditions of Methylenecyclo-propanes with Carbon Monoxide. *Angew. Chem., Int. Ed.* **2005**, *44*, 7881. (c) de Meijere, A.; Kurahashi, T. [5 + 1] Cycloaddition of (Cyclopropylmethylene) Cyclo-propanes and Other Vinylcyclo-propanes with Carbon Monoxide Catalyzed by Octacarbonyldi-cobalt. *Synlett* **2005**, 2619.
- (6) For Co(I)-mediated C-C bond cleavage, see: (a) Liebeskind, L. S.; Baysdon, S. L.; Goedken, V.; Chidambaram, R. Phthaoylcobalt Complexes in Synthesis. Ligand Modifications Leading to a Practical Naphthoquinone Synthesis. *Organometallics* **1986**, *5*, 1086. (b) Huffman, M. A.; Liebeskind, L. S. Insertion of ( $\eta^5$ -Indenyl)cobalt(I) into Cyclobutenones: The First Synthesis of Phenols from Isolated Vinylketene Complexes. *J. Am. Chem. Soc.* **1990**, *112*, 8617. (c) Huffman, M. A.; Liebeskind, L. S.; Pennington, W. T. Reaction of Cyclobutenones with Low-Valent Metal Reagents To Form  $\eta^4$ - and  $\eta^5$ -Vinylketene Complexes. Reaction of  $\eta^4$ -Vinylketene Complexes with Alkynes To Form Phenols. *Organometallics* **1992**, *11*, 255.
- (7) (a) Bourne-Branchu, Y.; Gosmini, C.; Danoun, G. Cobalt-Catalyzed Esterification of Amides. *Chem. - Eur. J.* **2017**, *23*, 10043. (b) Dorval, C.; Dubois, E.; Bourne-Branchu, Y.; Gosmini, C.; Danoun, G. Sequential Organozinc Formation and Negishi Cross-Coupling of Amides Catalysed by Cobalt Salt. *Adv. Synth. Catal.* **2019**, *361*, 1777.
- (8) For selected examples of Rh-catalyzed catalyzed decarbonylation of amide, see: (a) Meng, G.; Szostak, M. Rhodium-Catalyzed C-H Bond Functionalization with Amides by Double C-H/C-N Bond Activation. *Org. Lett.* **2016**, *18*, 796. For selected examples of Palladium-catalyzed decarbonylation of amide, see: (b) Meng, G.; Szostak, M. General Olefin Synthesis by the Palladium-Catalyzed Heck Reaction of Amides: Sterically Controlled Chemoselective N-C Activation. *Angew. Chem., Int. Ed.* **2015**, *54*, 14518. (c) Meng, Y. Y.; Si, X. J.; Song, Y. Y.; Zhou, H. M.; Xu, F. Palladium-Catalyzed Decarbonylative Annulation of Phthalimides with Arynes: Direct Construction of Phenanthridinones. *Chem. Commun.* **2019**, *55*, 9507. (d) Shi, S.; Szostak, M. Decarbonylative Borylation of Amides by Palladium Catalysis. *ACS Omega* **2019**, *4*, 4901. For selected examples of Ni-catalyzed decarbonylation of amides, see: (e) Kajita, Y.; Matsubara, S.; Kurahashi, T. Nickel-Catalyzed Decarbonylative Addition of Phthalimides to Alkynes. *J. Am. Chem. Soc.* **2008**, *130*, 6058. (f) Shiba, T.; Kurahashi, T.; Matsubara, S. Nickel-Catalyzed Decarbonylative Alkylenation of Phthalimides with Trimethylsilyl-Substituted Alkynes. *J. Am. Chem. Soc.* **2013**, *135*, 13636. (g) Mi, P.; Liao, P.; Tu, T.; Bi, X. Decarbonylative C-C Bond-Forming Reactions of Saccharins by Nickel Catalysis: Homocoupling and Cycloaddition. *Chem. - Eur. J.* **2015**, *21*, 5332. (h) Hu, F.; Lalancette, R.; Szostak, M. Structural Characterization of N-Alkylated Twisted Amides: Consequences for Amide Bond Resonance and N-C Cleavage. *Angew. Chem., Int. Ed.* **2016**, *55*, 5062. (i) Hu, J.; Zhao, Y.; Liu, J.; Zhang, Y.; Shi, Z. Nickel-Catalyzed Decarbonylative Borylation of Amides: Evidence for Acyl C-N Bond Activation. *Angew. Chem., Int. Ed.* **2016**, *55*, 8718. (j) Liu, X.; Yue, H.; Jia, J.; Guo, L.; Rueping, M. Synthesis of Amidines from Amides Using a Nickel-Catalyzed Decarbonylative Amination through CO Extrusion Intramolecular Recombination Fragment Coupling. *Chem. - Eur. J.* **2017**, *23*, 11771. (k) Yue, H.; Guo, L.; Liao, H. H.; Cai, Y.; Zhu, C.; Rueping, M. Catalytic Ester and Amide to Amine Interconversion: Nickel-Catalyzed Decarbonylative Amination of Esters and Amides by C-O and C-C Bond Activation. *Angew. Chem., Int. Ed.* **2017**, *56*, 4282. (l) Morioka, T.; Nakatani, S.; Sakamoto, Y.; Kodama, T.; Ogoshi, S.; Chatani, N.; Tobisu, M. Nickel-Catalyzed Decarbonylation of N-Acylated N-Heteroarenes. *Chem. Sci.* **2019**, *10*, 6666. Relevant review on terminology of amides, see: (m) Liu, C.; Szostak, M. Twisted Amides: From Obscurity to Broadly Useful Transition-Metal-Catalyzed Reactions by N-C Amide Bond Activation. *Chem. - Eur. J.* **2017**, *23*, 7157. Relevant review on decarbonylative reactions of amides, see: (n) Liu, C.; Szostak, M. Decarbonylative cross-coupling of amides. *Org. Biomol. Chem.* **2018**, *16*, 7998.

- (9) (a) Komatsuda, M.; Kato, H.; Muto, K.; Yamaguchi, J. Pd-Catalyzed Dearomatic Three-Component Reaction of Bromoarenes with Diazo Compounds and Allylborates. *ACS Catal.* **2019**, *9*, 8991. (b) Pan, N.; Ling, J.; Zapata, R.; Pulicani, J.-P.; Grimaud, L.; Vitale, M. R. Electrochemical TEMPO-Catalyzed Multicomponent C(sp<sup>3</sup>)-H α-Carbamoylation of Free Cyclic Secondary Amines. *Green Chem.* **2019**, *21*, 6194. (c) van Vliet, K. M.; Polak, L. H.; Siegler, M. A.; van der Vlugt, J. I.; Guerra, C. F.; de Bruin, B. Efficient Copper-Catalyzed Multicomponent Synthesis of N-Acyl Amidines via Acyl Nitrenes. *J. Am. Chem. Soc.* **2019**, *141*, 15240. (d) Yang, J.; Ji, D.-W.; Hu, Y.-C.; Min, X.-T.; Zhou, X.; Chen, Q.-A. Cobalt-Catalyzed Hydroxymethylation of Terpenes with Formaldehyde and Arenes. *Chem. Sci.* **2019**, *10*, 9560. (e) Yu, S.; Wu, J.; Lan, H.; Gao, L.; Qian, H.; Fan, K.; Yin, Z. Palladium and Bronsted Acid Co-Catalyzed Biginelli-Like Multicomponent Reactions via in Situ-Generated Cyclic Enol Ether: Access to Spirofuran-Hydropyrimidinones. *Org. Lett.* **2020**, *22*, 102.
- (10) (a) Qiu, S.; Zhai, S.; Wang, H.; Tao, C.; Zhao, H.; Zhai, H. Efficient Synthesis of Phthalimides via Cobalt-Catalyzed C(sp<sup>2</sup>)-H Carbonylation of Benzoyl Hydrazides with Carbon Monoxide. *Adv. Synth. Catal.* **2018**, *360*, 3271. (b) Grigorjeva, L.; Daugulis, O. Cobalt-Catalyzed Direct Carbonylation of Aminoquinoline Benzamides. *Org. Lett.* **2014**, *16*, 4688.
- (11) (a) Gallagher, A. G.; Tian, H.; Torres-Herrera, O. A.; Yin, S.; Xie, A.; Lange, D. M.; Wilson, J. K.; Mueller, L. G.; Gau, M. R.; Carroll, P. J.; Martinez-Solorio, D. Access to Highly Functionalized Cyclopentenones via Diastereo-selective Pauson-Khand Reaction of Siloxy-Tethered 1,7-Enynes. *Org. Lett.* **2019**, *21*, 8646. (b) Garçon, M.; Cabré, A.; Verdaguer, X.; Riera, A. Synthesis, Coordination Study, and Catalytic Pauson-Khand Reactions of Quinoxp\*(Co)<sub>4</sub>-M-Alkyne Dicobalt Complexes. *Organometallics* **2017**, *36*, 1056. (c) Orgue, S.; Leon, T.; Riera, A.; Verdaguer, X. Asymmetric Intermolecular Cobalt-Catalyzed Pauson-Khand Reaction Using a Pstereogenic Bis-Phosphane. *Org. Lett.* **2015**, *17*, 250.
- (12) (a) Wang, M.; Hu, Y.; Jiang, Z.; Shen, H. C.; Sun, X. Divergent Copper-Mediated Dimerization and Hydroxylation of Benzamides Involving C-H Bond Functionalization. *Org. Biomol. Chem.* **2016**, *14*, 4239. (b) Du, C.; Li, P.-X.; Zhu, X.; Han, J.-N.; Niu, J.-L.; Song, M.-P. Cobalt-Catalyzed Oxidative C-H/N-H Cross-Coupling: Selective and Facile Access to Triarylamines. *ACS Catal.* **2017**, *7*, 2810.
- (13) Yuan, Y. C.; Kamaraj, R.; Bruneau, C.; Labasque, T.; Roisnel, T.; Gramage-Doria, R. Unmasking Amides: Ruthenium-Catalyzed Protodecarbonylation of N-Substituted Phthalimide Derivatives. *Org. Lett.* **2017**, *19*, 6404.
- (14) (a) Mori, T.; Ito, T.; Liu, S.; Ando, H.; Sakamoto, S.; Yamaguchi, Y.; Tokunaga, E.; Shibata, N.; Handa, H.; Hakoshima, T. Structural Basis of Thalidomide Enantiomer Binding to Cereblon. *Sci. Rep.* **2018**, *8*, 1294. (b) Wójcik, P.; Trzeciak, A. M. The Aminocarbonylation of 1,2-Diodoarenes with Primary and Secondary Amines Catalyzed by Palladium Complexes with Imidazole Ligands. *Appl. Catal., A* **2018**, *560*, 73.
- (15) (a) Lohbeck, J.; Miller, A. K. Practical Synthesis of a Phthalimide-Based Cereblon Ligand to Enable Protac Development. *Bioorg. Med. Chem. Lett.* **2016**, *26*, 5260. (b) Okada-Junior, C. Y.; Monteiro, G. C.; Aguiar, A. C. C.; Batista, V. S.; de Souza, J. O.; Souza, G. E.; Bueno, R. V.; Oliva, G.; Nascimento-Junior, N. M.; Guido, R. V. C.; Bolzani, V. S. Phthalimide Derivatives with Bioactivity Against Plasmodium Falciparum: Synthesis, Evaluation, and Computational Studies Involving bc1 Cytochrome Inhibition. *ACS Omega* **2018**, *3*, 9424.
- (16) (a) Okamoto, H.; Konishi, H.; Satake, K. Fluorescence Response of 3-Trifluoroacetylaminophthalimide to a Li<sup>+</sup>-I<sup>-</sup> Ion Pair Induced by 254 nm Photolysis in Acetonitrile. *Chem. Commun.* **2012**, *48*, 2346. (b) Huang, J.; Chen, B.; Zhou, B.; Han, Y. A Novel ESIPT-Based Fluorescent Chemodosimeter for Hg<sup>2+</sup> Detection and Its Application in Live-Cell Imaging. *New J. Chem.* **2018**, *42*, 1181. (c) Mayer, M.; Takegami, S.; Neumeier, M.; Rink, S.; Jacobi von Wangenheim, A.; Schulte, S.; Vollmer, M.; Griesbeck, A. G.; Duerkop, A.; Baeumner, A. J. Electrochemiluminescence Bioassays with a Water-Soluble Luminol Derivative Can Outperform Fluorescence Assays. *Angew. Chem., Int. Ed.* **2018**, *57*, 408. Phthalimides are activated amides by two carbonyl groups, thus expected to be much more reactive than N-acyl amides, relevant references see: (d) Meng, G.; Shi, S.; Lalancette, R.; Szostak, R.; Szostak, M. Reversible Twisting of Primary Amides via Ground State N-C(O) Destabilization: Highly Twisted Rotationally Inverted Acyclic Amides. *J. Am. Chem. Soc.* **2018**, *140*, 727.
- (17) Kalsi, D.; Dutta, S.; Barsu, N.; Rueping, M.; Sundararaju, B. Room-Temperature C-H Bond Functionalization by Merging Cobalt and Photoredox Catalysis. *ACS Catal.* **2018**, *8*, 8115.
- (18) Zhang, T. Y.; Lin, J. B.; Li, Q. Z.; Kang, J. C.; Pan, J. L.; Hou, S. H.; Chen, C.; Zhang, S. Y. Copper-Catalyzed Selective Ortho-C-H/N-H Annulation of Benzamides with Arynes: Synthesis of Phenanthridinone Alkaloids. *Org. Lett.* **2017**, *19*, 1764.
- (19) Hickey, D. P.; Sandford, C.; Rhodes, Z.; Gensch, T.; Fries, L. R.; Sigman, M. S.; Minteer, S. D. Investigating the Role of Ligand Electronics on Stabilizing Electrocatalytically Relevant Low-Valent Co(I) Intermediates. *J. Am. Chem. Soc.* **2019**, *141*, 1382.