

ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: D. Ji, G. He, W. Zhang, C. Zhao, Y. Hu and Q. Chen, *Chem. Commun.*, 2020, DOI: 10.1039/D0CC02697B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

Nickel-catalyzed allyl-allyl coupling reactions between 1,3-dienes and allylboronates

Received 00th January 20xx,
Accepted 00th January 20xxDing-Wei Ji,^{a,b} Gu-Cheng He,^a Wei-Song Zhang,^a Chao-Yang Zhao,^a Yan-Cheng Hu,^a Qing-An Chen^{*a}


DOI: 10.1039/x0xx00000x

A regioselective allyl-allyl coupling reaction between 1,3-dienes and allylboronates has been demonstrated under nickel catalysis. Salient features of this method include the earth-abundant metal catalyst, excellent regioselectivity and good functional group tolerance. Notably, even congested allyl substrates can also be applied to this protocol, thus allowing for the rapid preparation of a series of valuable 1,5-dienes.

Owing to the diverse reactivity of the allyl and vinyl groups, 1,5-diene fragments and their derivatives often serve as useful building blocks in organic synthesis.¹ Besides, 1,5-diene unit itself is widely found in biologically active molecules and natural products.² Therefore, their synthesis has attracted considerable attention over the past decades and some efficient strategies have been established.³ Among them, transition metal-catalyzed cross-coupling between allyl electrophiles and allyl nucleophiles represents one of the simplest and most straightforward tools.⁴ However, two inherent challenges limit the applications of this classic strategy (Scheme 1a). First, the requirement for pre-installation of a leaving group onto the starting allyl electrophiles may diminish the synthetic economy. Second, several difficulties such as the poor regioselective control of products or the competing formation of side product 1,3-dienes derived from β -hydride elimination may arise when utilizing internal allyl electrophiles as the coupling partners.^{4i,4o} In this context, developing a general and regioselective allyl-allyl coupling reaction to access 1,5-dienes from non-activated precursors is still highly demanded.

As 1,3-dienes are easily accessible and versatile synthons in organic synthesis, the employment of 1,3-dienes as greener allyl precursors via the catalytic hydrofunctionalization is becoming an attractive way for the buildup of molecular

complexity.^{5,6} Despite the fact that reductive allyl-allyl cross-coupling of allenes or alkynes have been developed by Hoveyda,⁷ Tsuji,⁸ Zhang⁹ and our group,¹⁰ the direct utilization of 1,3-dienes in allyl-allyl coupling for the synthesis of 1,5-dienes was rarely realized. Very recently, Meek and co-workers reported a seminal work on the rhodium-catalyzed hydroallylation of 1,3-dienes with allyltrifluoroborates.¹¹ Due to undesired alkene isomerization, a mixture of 1,5- and 1,4-dienes were obtained in most cases (Scheme 1b). In light of excellent performance of nickel catalyst in 1,3-diene hydrofunctionalization,¹² we envisioned that the relatively cheap and earth-abundant nickel could also serve as an attractive alternative to noble rhodium catalyst in hydroallylation of 1,3-dienes. As an extension of our studies on catalytic functionalization of dienes,¹³ herein we report this regioselective reductive allyl-allyl cross-coupling between 1,4-dienes and allylboronates under nickel-hydride catalysis (Scheme 1c).




Scheme 1. Transition metal-catalyzed allyl-allyl coupling reactions

^a Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian116023, P. R. China, E-mail: qachen@dicp.ac.cn, Web: http://www.lbc.dicp.ac.cn^b University of Chinese Academy of Sciences, Beijing 100049, P. R. China
Electronic Supplementary Information (ESI) available: See
DOI: 10.1039/x0xx00000x

We initiated our research with 1-phenyl-1,3-butadiene **1a** and allyl-B(pin) **2a** as the model substrates (Table 1). Pleasingly, the expected 1,5-diene **3a** was observed as sole product in the presence of Ni(COD)₂ (5 mol%) and PPh₃ (10 mol%) in MeOH at 80 °C (entry 1). The evaluation of solvent suggested that the yield of **3a** could be improved to 58% when the reaction was performed in ethanol (entries 2-5). After examining various ligands, we obtained an increased yield of **3a** with P(^{*n*}Bu)₃ (entries 6-11). While lowering or raising temperature resulted in decreased yields (entries 12-13), increasing the reaction concentration successfully led to **3a** in an excellent yield (93%, entry 14).

Table 1. Optimization for hydroallylation of 1,3-diene




entry	ligand (x mol%)	solvent	yield (%) ^b
1	PPh ₃ (10)	MeOH	50
2	PPh ₃ (10)	EtOH	58
3	PPh ₃ (10)	^{<i>n</i>} PrOH	51
4	PPh ₃ (10)	^{<i>i</i>} PrOH	55
5	PPh ₃ (10)	^{<i>t</i>} BuOH	45
6	PCy ₃ (10)	EtOH	22
7	P(^{<i>n</i>} Bu) ₃ (10)	EtOH	trace
8	P(^{<i>n</i>} Bu) ₃ (10)	EtOH	83
9	dppp (5)	EtOH	2
10	dppb (5)	EtOH	50
11	dppf (5)	EtOH	5
12 ^c	P(^{<i>n</i>} Bu) ₃ (10)	EtOH	42
13 ^d	P(^{<i>n</i>} Bu) ₃ (10)	EtOH	74
14 ^e	P(^{<i>n</i>} Bu) ₃ (10)	EtOH	93 (85) ^g
15 ^f	P(^{<i>n</i>} Bu) ₃ (10)	EtOH	75

^aReaction conditions: **1a** (0.20 mmol), **1b** (0.40 mmol), Ni(COD)₂ (5 mol%), Ligand (5-10 mol%), solvent (0.4 M), 80 °C, 18 h. ^bDetermined by GC-FID with 1,3,5-trimethoxybenzene as the internal standard. ^c60 °C. ^d100 °C. ^eEtOH (0.8 M), ^fEtOH (0.2 M). ^gIsolated yield.

With the optimal reaction conditions in hand, we shifted our attention to explore the substrate scope of this nickel-catalyzed allyl-allyl coupling reaction. As shown in Table 2, we generally obtained various 1,5-dienes in good yields and selectivities. The 1,3-dienes with alkyl substituents on phenyl ring such as methyl, ethyl and butyl all gave desirable products in good to excellent yields (**3b-3d**, **3j**, **3k**). The strong electron-donating substrates (e.g. OMe, NMe₂) were comparatively sluggish in this reaction (**3e** and **3f**). A fluoro group was also well tolerated, delivering the 1,5-diene **3g** in 80% yield. Electron-withdrawing Ph and CF₃ groups exhibited higher reactivity and offered good yields of 1,5-dienes even at lower temperature (**3h**, **3i** and **3l**). Notably, *ortho*-substituted substrates **1m** and **1n** also proceeded smoothly, regardless of their steric hindrance, generating the corresponding products in 83% and 85% yield, respectively. On replacing the phenyl group with 1-naphthalenyl group, the substrate was

transformed into product **3o** in an excellent yield. Heterocyclic substrates **3p-3r** were also compatible in current protocol and afforded the products in 21-60% yields.

Table 2. Hydroallylation of 1,3-diene



Reaction conditions: **1** (0.20 mmol), **2a** (0.40 mmol), Ni(COD)₂ (5 mol%), P^{*n*}Bu₃ (10 mol%), EtOH (0.8 M), 80 °C, isolated yields were given. The rr values refer to the ratio of **3** to 1,4 adducts and determined by ¹H NMR analysis. Unless noted, the rr is more than 20:1. ^aAccompanied by small amount of inseparable diene **1**, the yield of product has been adjusted accordingly. ^b70 °C. ^c60 °C.


The transition-metal catalyzed allyl-allyl couplings with congested allylboronates is a challenging task and very limited works have been successfully realized.^{4j,4o} Satisfactorily, although linear substrate **2b** was unreactive under current reaction conditions, the prenylative product **3s** could be obtained in an acceptable yield with branched prenyl-B(pin) (Table 3). Moreover, geranyl-B(pin) **2d** also reacted well with **1a** and gave the product **3t** in 42% yield. These results complement the Rh-catalyzed hydroallylation method for the synthesis of 1,5-dienes.¹¹

Table 3. The scope of allylborons


2	product	yield (%)
		< 5%
		43%
		42%
		Z/E mixture, 1/1.6 ^a

Reaction conditions: **1a** (0.20 mmol), **2** (0.40 mmol), Ni(COD)₂ (5 mol%), P^tBu₃ (10 mol%), EtOH (0.8 M), 80 °C, 18 h, isolated yields were given, unless noted, the *rr* > 20:1. ^adetermined by GC-FID.


To further verify the synthetic utility of this protocol, a scale-up reaction (5.0 mmol) was performed, delivering 1,5-diene product **3a** in 88% yield (762 mg, Scheme 2). Besides, further synthetic transformation of **3a** with phenyl acrylate under cross olefin-metathesis conditions furnished the unsaturated ester **4** in 37% yield. Moreover, the hydroboration/oxidation reactions of **3a** could proceed efficiently to give the alcohol **5** in a good yield (82%).

**Scheme 2.** Scale-up synthesis and transformations of 1,5-diene **3a**

To gain some insights into the mechanism, a deuterium-labeling experiment with 1-phenyl-1,3-butadiene **1a** and allyl-B(pin) **2a** was conducted with EtOH-*d*₆. As depicted in Scheme 3a, while 0.37 deuterium was incorporated into the methyl group of 1,5-diene product, 0.64 deuterium was also exchanged into terminal CH₂ of the recovered diene **1a**. This observation supports a reversible process of the migratory insertion between the 1,4-diene and the Ni-hydride. In the absence of **2a**, hydroalkoxylation product **6** could also be obtained in 21% NMR yield under the standard reaction condition. However, the reaction between **6** and **2a** under standard condition could not deliver 1,5-diene **3a** (Scheme 3b), thus excluding the possibility that **6** is an intermediate of the process.

**Scheme 3.** Deuterium-Labeling and Control Experiments

While further studies are warranted, we tentatively proposed the following mechanism on the basis of literature and our own observations (Figure 1).^{11,12f} First, the oxidative addition of the ethanol with a nickel(0) precursor **A** gives Ni(II)-hydride species **B**, which then undergoes migratory insertion with 1,3-diene **1** to afford allyl-Ni(II) intermediate **C** or **C'** in a reversible pathway. Through the transmetalation between allyl-Ni(II) intermediate and allyl-B(pin) **2**, bis(allyl)Ni species **D** is produced. Finally, reductive elimination of **D** leads to terminal 1,5-diene product **3** and regenerates the nickel(0). It should be noted that the existence of allyl-Ni(II) intermediate **C** / **C'** were supported by the observation of **6** via a reductive elimination step. Therefore, current mechanism was preferred over the reported transmetalation-alkene insertion pathway in previous Rh-catalyzed reactions.¹¹

**Figure 1.** Proposed mechanism for Ni-catalyzed hydroallylation

In summary, a nickel-catalyzed allyl-allyl cross-coupling reaction between 1,3-dienes and allyl-B(pin) was achieved via a hydroallylative protocol. This strategy exhibited good functional group tolerance, exclusive regioselectivity and high reactivity, thus complementing a facile route to access 1,5-dienes. Further studies on expanding the scope of unsaturated substrates, detailed mechanism as well as the asymmetric version are now in progress in our laboratory.

Conflicts of interest

There are no conflicts to declare.


Acknowledgements

Financial support from Dalian Institute of Chemical Physics (DICPI201902), the National Natural Science Foundation of China (21702204), Liaoning Revitalization Talents Program (XLYC1807181), and the "Thousand Youth Talents Plan" is acknowledged.

Notes and references

- (a) H.-Y. Lee, *Acc. Chem. Res.*, 2015, **48**, 2308; (b) J. Adrian, S. Roth and C. B. W. Stark, *ChemCatChem*, 2016, **8**, 1679; (c) P. K. Dornan, D. Lee and R. H. Grubbs, *J. Am. Chem. Soc.*, 2016, **138**, 6372; (d) Y. R. Gao, D. Y. Wang and Y. Q. Wang, *Org. Lett.*, 2017, **19**, 3516; (e) B. Schmidt, M. H. Petersen and D. Braun, *J. Org. Chem.*, 2018, **83**, 1627.
- (a) E. Breitmaier, *Terpenes: Flavors, Fragrances, Pharmaca, Pheromones*, Wiley-VCH, Weinheim, Germany, 2006; (b) L. Chuang, C. H. Wen, Y. R. Lee, Y. L. Lin, L. R. Hsu, S. Y. Wang and F. H. Chu, *J. Nat. Prod.*, 2018, **81**, 1162; (c) A. C. Huang, Y. J. Hong, A. D. Bond, D. J. Tantillo and A. Osbourn, *Angew. Chem. Int. Ed.*, 2018, **57**, 1291.
- (a) Y. Sumida, S. Hayashi, K. Hirano, H. Yorimitsu and K. Oshima, *Org. Lett.*, 2008, **10**, 1629; (b) F. Ding, R. William, F. Wang and X.-W. Liu, *Chem. Commun.*, 2012, **48**, 8709; (c) J. Y. Hamilton, D. Sarlah and E. M. Carreira, *J. Am. Chem. Soc.*, 2014, **136**, 3006; (d) Q. Yuan, K. Yao, D. Liu and W. Zhang, *Chem. Commun.*, 2015, **51**, 11834; (e) M. Ellwart, I. S. Makarov, F. Achraimer, H. Zipse and P. Knochel, *Angew. Chem. Int. Ed.*, 2016, **55**, 10502; (f) H. Wang, S. Qiu, S. Wang and H. Zhai, *ACS Catal.*, 2018, **8**, 11960.
- (a) P. Srihari, A. P. Singh, A. K. Basak and J. S. Yadav, *Tetrahedron Lett.*, 2007, **48**, 5999; (b) P. Zhang, L. A. Brozek and J. P. Morken, *J. Am. Chem. Soc.*, 2010, **132**, 10686; (c) L. A. Brozek, M. J. Ardolino and J. P. Morken, *J. Am. Chem. Soc.*, 2011, **133**, 16778; (d) T. Chatterjee, R. Dey and B. C. Ranu, *New J. Chem.*, 2011, **35**, 1103; (e) A. Jimenez-Aquino, E. F. Flegeau, U. Schneider and S. Kobayashi, *Chem. Commun.*, 2011, **47**, 9456; (f) P. Zhang, H. Le, R. E. Kyne and J. P. Morken, *J. Am. Chem. Soc.*, 2011, **133**, 9716; (g) M. B. Li, Y. Wang and S. K. Tian, *Angew. Chem. Int. Ed.*, 2012, **51**, 2968; (h) V. Hornillos, M. Perez, M. Fananas-Mastral and B. L. Feringa, *J. Am. Chem. Soc.*, 2013, **135**, 2140; (i) H. Le, R. E. Kyne, L. A. Brozek and J. P. Morken, *Org. Lett.*, 2013, **15**, 1432; (j) M. J. Ardolino and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 7092; (k) J. Y. Hamilton, N. Hauser, D. Sarlah and E. M. Carreira, *Angew. Chem. Int. Ed.*, 2014, **53**, 10759; (l) H. Le, A. Batten and J. P. Morken, *Org. Lett.*, 2014, **16**, 2096; (m) Y. Yasuda, H. Ohmiya and M. Sawamura, *Angew. Chem. Int. Ed.*, 2016, **55**, 10816; (n) X. B. Wang, X. M. Wang, Z. B. Han, Z. Wang and K. L. Ding, *Angew. Chem. Int. Ed.*, 2017, **56**, 1116; (o) Y. Zheng, B.-B. Yue, K. Wei and Y.-R. Yang, *Org. Lett.*, 2018, **20**, 8035.
- (a) N. Herrmann, D. Vogelsang, A. Behr and T. Seidensticker, *ChemCatChem*, 2018, **10**, 5342; (b) M. Holmes, L. A. Schwartz and M. J. Krische, *Chem. Rev.*, 2018, **118**, 6026; (c) Y. Xiong, Y. W. Sun and G. Z. Zhang, *Tetrahedron Lett.*, 2018, **59**, 347; (d) N. J. Adamson and S. J. Malcolmson, *ACS Catal.*, 2020, **10**, 1060; (e) X. Wu and L. Z. Gong, *Synthesis*, 2019, **51**, 122; (f) G. J. P. Perry, T. Jia and D. J. Procter, *ACS Catal.*, 2020, **10**, 1485. View Article Online
DOI: 10.1039/D0CC02697B
- (a) N. J. Adamson, E. Hull and S. J. Malcolmson, *J. Am. Chem. Soc.*, 2017, **139**, 7180; (b) Y. Y. Gui, N. Hu, X. W. Chen, L. L. Liao, T. Ju, J. H. Ye, Z. Zhang, J. Li and D. G. Yu, *J. Am. Chem. Soc.*, 2017, **139**, 17011; (c) J. S. Marcum, C. C. Roberts, R. S. Manan, T. N. Cervarich and S. J. Meek, *J. Am. Chem. Soc.*, 2017, **139**, 15580; (d) X. H. Yang and V. M. Dong, *J. Am. Chem. Soc.*, 2017, **139**, 1774; (e) X. H. Yang, A. Lu and V. M. Dong, *J. Am. Chem. Soc.*, 2017, **139**, 14049; (f) N. J. Adamson, K. C. E. Wilbur and S. J. Malcolmson, *J. Am. Chem. Soc.*, 2018, **140**, 2761; (g) S. Z. Nie, R. T. Davison and V. M. Dong, *J. Am. Chem. Soc.*, 2018, **140**, 16450; (h) X. H. Yang, R. T. Davison and V. M. Dong, *J. Am. Chem. Soc.*, 2018, **140**, 10443; (i) X. W. Chen, L. Zhu, Y. Y. Gui, K. Jing, Y. X. Jiang, Z. Y. Bo, Y. Lan, J. Li and D. G. Yu, *J. Am. Chem. Soc.*, 2019, **141**, 18825; (j) G. Wang, L. Gao, H. Chen, X. Liu, J. Cao, S. Chen, X. Cheng and S. Li, *Angew. Chem. Int. Ed.*, 2019, **58**, 1694; (k) X. H. Yang, R. T. Davison, S. Z. Nie, F. A. Cruz, T. M. McGinnis and V. M. Dong, *J. Am. Chem. Soc.*, 2019, **141**, 3006.
- (a) F. K. Meng, K. P. McGrath and A. H. Hoveyda, *Nature*, 2014, **513**, 367; (b) Y. Sun, Y. Zhou, Y. Shi, J. Del Pozo, S. Torker and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2019, **141**, 12087.
- K. Semba, N. Bessho, T. Fujihara, J. Terao and Y. Tsuji, *Angew. Chem. Int. Ed.*, 2014, **53**, 9007.
- G. Xu, B. Fu, H. Zhao, Y. Li, G. Zhang, Y. Wang, T. Xiong and Q. Zhang, *Chem. Sci.*, 2019, **10**, 1802.
- D.-W. Ji, Y.-C. Hu, H. Zheng, C.-Y. Zhao, Q.-A. Chen and V. M. Dong, *Chem. Sci.*, 2019, **10**, 6311.
- J. S. Marcum, T. N. Cervarich, R. S. Manan, C. C. Roberts and S. J. Meek, *ACS Catal.*, 2019, **9**, 5881.
- (a) J. Pawlas, Y. Nakao, M. Kawatsura and J. F. Hartwig, *J. Am. Chem. Soc.*, 2002, **124**, 3669; (b) N. Saito, A. Kobayashi and Y. Sato, *Angew. Chem. Int. Ed.*, 2012, **51**, 1228; (c) A. Koepfer, B. Sam, B. Breit and M. J. Krische, *Chem. Sci.*, 2013, **4**, 1876; (d) X. Lian, W. Chen, L. Dang, Y. Li and C. Y. Ho, *Angew. Chem. Int. Ed.*, 2017, **56**, 9048; (e) L. Cheng, M. M. Li, L. J. Xiao, J. H. Xie and Q. L. Zhou, *J. Am. Chem. Soc.*, 2018, **140**, 11627; (f) L. J. Xiao, L. Cheng, W. M. Feng, M. L. Li, J. H. Xie and Q. L. Zhou, *Angew. Chem. Int. Ed.*, 2018, **57**, 461; (g) B. Chen, Y. Zhang, R. Wu, D. Fang, X. Chen, S. Wang, Y. Zhao, P. Hu, K.-Q. Zhao, B.-Q. Wang and P. Cao, *ACS Catal.*, 2019, **9**, 11788; (h) Y. G. Chen, B. Shuai, X. T. Xu, Y. Q. Li, Q. L. Yang, H. Qiu, K. Zhang, P. Fang and T. S. Mei, *J. Am. Chem. Soc.*, 2019, **141**, 3395; (i) J. Long, P. Wang, W. Wang, Y. Li and G. Yin, *iScience*, 2019, **22**, 369; (j) L. Lv, D. Zhu, Z. Qiu, J. Li and C.-J. Li, *ACS Catal.*, 2019, **9**, 9199; (k) X.-Y. Lv, C. Fan, L.-J. Xiao, J.-H. Xie and Q.-L. Zhou, *CCS Chem.*, 2019, **1**, 328; (l) G. Tran and C. Mazet, *Org. Lett.*, 2019, **21**, 9124; (m) G. Tran, W. Shao and C. Mazet, *J. Am. Chem. Soc.*, 2019, **141**, 14814; (n) Y.-L. Li, W.-D. Li, Z.-Y. Gu, J. Chen and J.-B. Xia, *ACS Catal.*, 2020, **10**, 1528.
- (a) Y. C. Hu, D. W. Ji, C. Y. Zhao, H. Zheng and Q. A. Chen, *Angew. Chem. Int. Ed.*, 2019, **58**, 5438; (b) J. Yang, D. W. Ji, Y. C. Hu, X. T. Min, X. G. Zhou and Q. A. Chen, *Chem. Sci.*, 2019, **10**, 9560; (c) Y. Li, Y. C. Hu, H. Zheng, D. W. Ji, Y. F. Cong and Q. A. Chen, *Eur. J. Org. Chem.*, 2019, 6510; (d) Y. Li, Y. C. Hu, D. W. Ji, W. S. Zhang, G. C. He, Y. F. Cong and Q. A. Chen, *Chin. J. Catal.*, 2020, DOI: 10.1016/S1872-2067(20)63575-6.

Graphical Abstracts (Table of Contents Entry)



A nickel-hydride catalysis has been developed to facilitate the allyl-allyl cross-coupling reactions between 1,3-dienes and allyl-B(pin) in excellent regioselectivity.