

Regioselective Hydroacylation of 1,3-Dienes by Cobalt Catalysis

Qing-An Chen, Daniel K. Kim, and Vy M. Dong*


Department of Chemistry, University of California, Irvine, 4403 Natural Sciences 1, Irvine, California 92697, United States

S Supporting Information


ABSTRACT: We describe a cobalt-catalyzed hydroacylation of 1,3-dienes with non-chelating aldehydes. Aromatic aldehydes provide 1,4-addition products as the major isomer, while aliphatic aldehydes favor 1,2-hydroacylation products. The kinetic profile supports an oxidative cyclization mechanism involving a cobaltacycle intermediate that undergoes transformation with high regio- and stereoselectivity.

A modern challenge in organic synthesis is the invention of methods that use catalysts derived from first-row transition metals.¹ A number of valuable olefin transformations, including hydrogenation,² hydroformylation,³ and hydrovinylation,⁴ have been achieved by cobalt catalysis. These breakthroughs highlight Co as an attractive and complementary alternative to Rh due to its relatively low cost and high abundance. Encouraged by this progress, we recently turned our attention to developing olefin hydroacylation^{5–14} by cobalt catalysis.¹⁵ Brookhart demonstrated the first and only previously known intermolecular Co-catalyzed hydroacylation (eq 1).^{15a,b} While promising, this


Brookhart's Work: C–H activation



This Work: Oxidative Cyclization




Krische and Ryu's Work: Hydrovinylation



strategy was limited to vinylsilanes, and aldehyde decarbonylation remained competitive. We imagined developing a Co(I)-catalyzed cross-coupling of various aldehydes and 1,3-dienes (eq 2). Our C–H bond functionalization would yield β,γ - and/or γ,δ -unsaturated ketones with high regioselectivity and excellent atom economy. Moreover, this proposed hydroacylation would afford regiocontrol distinct from current technologies, including the relevant Ru-catalyzed pathway developed independently by Krische^{11c} and Ryu^{11d} (eq 3).

Our proposal draws from a mechanism established by Hilt for the Co-catalyzed hydrovinylation of olefins to generate 1,4-dienes.⁴ As shown in Scheme 1, a Co(II) precatalyst can be used to generate the Co(I) catalyst.¹⁶ We hypothesize that Co(I) catalyst A would promote oxidative cyclization between aldehyde

Scheme 1. Proposed Cobalt-Catalyzed Hydroacylation of Dienes by Oxidative Cyclization



1 and a 1,3-diene 2.¹⁷ The resulting cobaltacycle B exists in two η^1 forms: seven-membered cobaltacycle C and five-membered cobaltacycle C'.^{4,18,19} Depending on the substrate and ligand, cobaltacycle C or C' would undergo transformation by Path I or II, respectively. A β -hydride elimination generates intermediate D or D', which can undergo reductive elimination to yield product 3 or 3' and regenerate catalyst A. In Path I, formation of the cobaltacycle intermediate requires a *cis*-olefin geometry; thus, we propose this pathway would lead to the *Z*-isomer of 3 with high levels of stereocontrol.

To test our hypothesis, we chose benzaldehyde 1a and isoprene 2a as model substrates. Control experiments confirm that these substrates are unreactive in the absence of either Co(II) salts or reducing agents.²⁰ We then examined the transformation using Co(II) (a catalyst precursor for hydrovinylation)⁴ with various ligands and additives to generate the requisite Co(I) catalyst. A survey of 20 commercially available phosphines reveals 1,3-bis(diphenylphosphino)propane (dppp) is the most promising ligand. With Co(II)-dppp, the desired product is observed in 11% yield by using Zn/ZnI₂, a known protocol for generating Co(I) (Table 1, entry 1).^{4,16} By applying In as the reductant instead, we observe improved reactivity (21%, entry 2). Through a further survey of additives, we find that a combination of In/InBr₃ gives the best result (57% yield, entries 3–5). Given the role of ZnI₂ in hydrovinylation,⁴ we assume InBr₃ similarly promotes the formation of cationic Co(I) species. In the absence of these Lewis acids, the yield is diminished (13% yield, entry 6). In all cases, we observe the β,γ -unsaturated ketone in preference to the γ,δ -unsaturated ketone (up to 19:1 selectivity). In accordance with our proposed mechanism, the

Received: January 9, 2014

Published: March 3, 2014

Table 1. Additives Effects for Hydroacylation^a

dppp = Ph₂P(CH₂)₃PPh₂

entry	additives	yield (%) ^b	3aa/3aa' ^c	Z/E (3aa) ^c
1	Zn/ZnI ₂	11	12:1	10:1
2	In/ZnI ₂	21	14:1	16:1
3	In/InI ₃	45	17:1	13:1
4	In/InBr ₃	57	19:1	11:1
5	In/InCl ₃	28	14:1	17:1
6	In/—	13	13:1	18:1

^a1a (0.20 mmol), 2a (0.60 mmol), Co(dppp)₂ (5 mol%), In or Zn (20 mol%), MX_n (5 mol%), DCE (1 mL), 60 °C, 24 h. ^bOverall yield of 3aa and 3aa', determined by ¹H NMR or GC-FID with dimethyl terephthalate (0.05 mmol) as internal standard. ^cDetermined by ¹H NMR or GC-FID.

Table 2. Ligand Effects of DPPP Analogues^a

Ar₂P(CH₂)₃PAr₂

L3: Ar = 4-FC₆H₄
 L4: Ar = 3,5-Me₂C₆H₃
 L1: Ar = 4-MeOC₆H₄
 L5: Ar = 3,5-(MeO)₂C₆H₃
 L2: Ar = 4-MeC₆H₄
 L6: Ar = 3,4-(MeO)₂C₆H₃

entry	catalyst	yield (%) ^b	3aa/3aa' ^c	Z/E (3aa) ^c
1	Co(dppp) ₂	57	19:1	11:1
2	Co(L1) ₂	59	17:1	17:1
3	Co(L2) ₂	63	19:1	13:1
4	Co(L3) ₂	50	14:1	15:1
5	Co(L4) ₂	83	>20:1	6:1
6	Co(L5) ₂	95	>20:1	3:1
7	Co(L6) ₂	83	>20:1	20:1
8 ^d	Co(L6) ₂	91 (87 ^e)	>20:1	19:1

^a1a (0.20 mmol), 2a (0.60 mmol), Co(II) (5 mol%), In (20 mol%), InBr₃ (5 mol%), DCE (1 mL), 60 °C, 24 h. ^bOverall yield of 3, determined by ¹H NMR or GC-FID with dimethyl terephthalate (0.05 mmol) as internal standard. ^cDetermined by ¹H NMR or GC-FID. ^dIn (10 mol%), DCE (0.5 mL). ^eIsolated yield.

stereochemistry of the resulting trisubstituted olefin is Z, as confirmed by 2D NOESY analysis.

To tune the catalyst, we prepared six analogues of dppp by varying the substitution pattern on the aryl groups (L1–L6, Table 2). From this study, we find that L6 (where Ar = 3,4-(MeO)₂C₆H₃) gives high yield, regioselectivity, and stereocontrol (87% isolated yield, >20:1 regioselectivity, and 19:1 Z/E, entry 8). With L6, the amount of In powder can be reduced to 10 mol% (entry 8). A trace amount of γ,δ -unsaturated ketone 4aa is observed as a minor product (3aa/4aa >20:1, entry 8). Our cobalt catalyst favors the 1,4-hydroacylation product 3aa over the 1,2-addition isomer 5aa which has been accessed with Ru(II) catalysis.^{11c,d} Thus, cobalt enables a rare type of hydroacylation that occurs across a conjugated π -system rather than a single π -bond.

With this protocol in hand, we explored the hydroacylation of diene 2a using 18 different aldehydes (Table 3). In general, good to high yields are obtained with various aromatic aldehydes 1 (60–97% yields, entries 1–13). In the case of electron-rich aryl aldehyde 1h, we observe a drop in stereoselectivity (2:1) due to

Table 3. Variation in the Aldehyde Scope^a

entry	R in 1	yield (%) ^b	3/3' ^c	Z/E (3) ^c
1	Ph (1a)	87	>20:1	19:1
2	4-BrC ₆ H ₄ (1b)	92	>20:1	>20:1
3	4-ClC ₆ H ₄ (1c)	96	>20:1	>20:1
4	4-FC ₆ H ₄ (1d)	60	>20:1	>20:1
5	4-CF ₃ C ₆ H ₄ (1e)	78	>20:1	>20:1
6	4-MeO ₂ CC ₆ H ₄ (1f)	88	>20:1	>20:1
7	4-MeC ₆ H ₄ (1g)	90	>20:1	6:1
8	4-MeOC ₆ H ₄ (1h)	97	>20:1	2:1
9 ^d	4-MeOC ₆ H ₄ (1h)	83	>20:1	11:1
10	3-BrC ₆ H ₄ (1i)	64	>20:1	>20:1
11	3-ClC ₆ H ₄ (1j)	89	>20:1	>20:1
12	2-furyl (1k)	94	>20:1	>20:1
13	2-naphthyl (1l)	94	>20:1	6:1
14	(E)-styryl (1m)	86	>20:1	6:1
15	Cy (1n)	17	2:1	n/a
16 ^e	Cy (1n)	82	1:8	n/a
17 ^e	n-C ₅ H ₁₁ (1o)	78	1:5	n/a
18 ^e	n-C ₆ H ₁₃ (1p)	83	1:4	n/a
19 ^e	n-C ₇ H ₁₅ (1q)	91	1:5	n/a
20 ^e	BnCH ₂ (1r)	74	1:4	n/a

^a1 (0.20 mmol), 2a (0.60 mmol), Co(L6)₂ (5 mol%), In (10 mol%), InBr₃ (5 mol%), DCE (0.5 mL), 60 °C, 20–24 h. ^bIsolated yield of all isomers. ^cDetermined by GC-FID or ¹H NMR; trace amount of product 4 was observed as minor isomer (3/4 >20:1) for aldehydes 1a–m. ^dDCE/toluene (1:1, 0.5 mL) was used solvent. ^e1 (0.20 mmol), 2a (0.60 mmol), Co(dcpe)₂ (5 mol%), In (20 mol%), InBr₃ (5 mol%), DCE/EtOAc (3:1, 0.5 mL), 50 °C, 20–24 h.

competitive olefin isomerization. The use of a mixed solvent (1:1 DCE/toluene) presumably inhibits isomerization and allows isolation of the desired product in good yield and higher stereoselectivity (11:1, entry 8 vs 9). This catalyst promotes hydroacylation of heteroaromatic aldehyde 1k (94% yield, entry 12). Moreover, the hydroacylation of isoprene 2a with α,β -unsaturated aldehyde 1m provides a conjugated ketone in 86% yield and >20:1 regioselectivity (entry 14).

With catalyst Co(L6)₂, aliphatic aldehyde 1n is less reactive and undergoes hydroacylation with lower regioselectivity (Table 3, entry 15). Toward addressing this challenge, we investigated a range of parameters and found promising reactivity with a more electron-rich phosphine ligand, 1,2-bis(dicyclohexylphosphino)ethane (dcpe).²⁰ The resulting ketone products are obtained in good yields with a dramatic switch in regiocontrol (entry 15 vs 16). Generally, the 1,2-hydroacylation products 3' are afforded as the major isomer with regioselectivities ranging from 4:1 up to 8:1 for aliphatic aldehydes 1n–r (entries 16–20). In these cases, we observe high C1-regioselectivity (C–C bond formation at 1-position of diene 1a) instead of C3-regioselectivity (eq 2 vs 3).

Next, we examined the scope of dienes (Table 4). For benzaldehyde, high yields (77–97%) and excellent regio- and stereoselectivities (>20:1 for both) are achieved for hydroacylation of 2-aryl-substituted butadienes, despite varying electronic and steric properties of substituents (entries 1–9). The catalyst loading can be reduced to 2 mol% for the coupling of benzaldehyde 1a and 2-phenylbutadiene (entry 2). The use of 2-cyclohexylbutadiene gives β,γ -unsaturated ketone 3aj with

Table 4. Variation of the Diene Partner^a



Entry	3	R in 3	Yield (%)
1		H	88 (3ab)
2		H	86 (3ab) ^b
3		4-Br	92 (3ac)
4		4-F	81 (3ad)
5		4-CF ₃	90 (3ae)
6		4-MeO	77 (3af)
7		4-Me	97 (3ag)
8		3-Cl	97 (3ah)
9		2-Cl	89 (3ai)
10		n/a	87 (3aj)
11		n/a	61 (3ak)
12		4-Br	78 (3bb)
13		4-F	73 (3db)
14		4-CF ₃	81 (3eb)
15		4-CO ₂ Me	88 (3fb)
16		4-Me	82 (3gb)
17		4-MeO	80 (3hb)

^a1 (0.30 mmol), 2 (0.20 mmol), Co(L6)I₂ (5 mol%), In (10 mol%), InBr₃ (5 mol%), DCE (0.5 mL), 60 °C, 16 h. Only one isomer was observed, except for 3aj (*E/Z* 6:1). The geometry of 3ab and 3ak was assigned by NOESY spectra. ^bCo(L6)I₂ (2 mol%).

moderate stereoselectivity (6:1) and good yield (87%, entry 10). Hydroacylation of a 1,2-disubstituted diene occurs with moderate yield (entry 11),²¹ while 2-phenylbutadiene **2b** couples well with a variety of aldehydes (entries 12–17). Throughout these studies, we observe no aldehyde decarbonylation, the byproduct expected for hydroacylations involving C–H bond activation of non-chelating aldehydes.

Finally, we report mechanistic studies that further support our proposed oxidative cyclization mechanism, in preference to a C–H activation pathway.^{15a,b} First, a kinetic isotope effect (KIE, k_H/k_D) of 2.7 is observed from competition experiments between benzaldehyde **1a** and *d*-**1a** (eq 8).²² If hydroacylation occurs through a traditional C–H activation mechanism, either oxidative addition of Co(I) to the aldehyde or insertion of the diene into the Co–H will be the turnover-limiting step.^{5d} However, we observe a zero-order dependence on both aldehyde and diene concentrations,^{20,23} thus disfavoring a C–H activation pathway. The KIE is consistent with β -hydride or reductive elimination as the turnover-limiting step in Path I (Scheme 1).²² Moreover, the zero-order dependence on both the aldehyde and the diene concentrations supports the possibility of metallacycle **C** or **D** as catalyst resting state (Scheme 1).

When the hydroacylation is performed with deuterio-benzaldehyde *d*-**1a**, the deuterium atom is incorporated at the



4-position of diene **2b** without any detectable deuterium at other positions (eq 9). This result suggests that the hydroacylation with aromatic aldehydes proceeds through a 1,4-addition pathway (Path I, Scheme 1). For hydroacylation with aliphatic aldehyde *d*-**1r**, the deuterium atom is incorporated completely at the β - and δ -position of products *d*-**3rb'** and *d*-**3rb**, respectively (eq 10). While further studies are warranted, our observations support a mechanistic proposal where Path I or II is favored (Scheme 1), depending on the properties of both the ligand and the substrate.

In contrast to the traditional mechanism of hydroacylation catalyzed by Rh(I) or Co(I), we propose an oxidative cyclization mechanism that avoids decarbonylation. Our catalyst promotes C1-regioselective hydroacylation of dienes and can be tuned to favor either 1,4- or 1,2-hydroacylation. Through the 1,4-hydroacylation pathway, we achieve the stereoselective synthesis of trisubstituted olefins, which are key building blocks that are challenging to access.²⁴ Ongoing efforts will focus on studying other catalysts for achieving higher selectivities and greater scope. Our study contributes to the emerging strategies available for hydroacylation via non-precious-metal catalysis.^{13,15}

■ ASSOCIATED CONTENT

📄 Supporting Information

Experimental details and kinetic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

✉ Corresponding Author

vy.dong@uci.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Funding was provided by UC Irvine and the National Institutes of Health (GM105938). V.M.D. is grateful to Eli Lilly for a Grantee Award.

■ REFERENCES

- (1) *Catalysis without Precious Metals*; Bullock, R. M., Ed.; Wiley-VCH: Weinheim, 2010.
- (2) For recent works on Co-catalyzed hydrogenations, see: (a) Monfette, S.; Turner, Z. R.; Semproni, S. P.; Chirik, P. J. *J. Am. Chem. Soc.* **2012**, *134*, 4561. (b) Zhang, G. Q.; Scott, B. L.; Hanson, S. K. *Angew. Chem., Int. Ed.* **2012**, *51*, 12102. (c) Friedfeld, M. R.; Shevlin, M.; Hoyt, J. M.; Krska, S. W.; Tudge, M. T.; Chirik, P. J. *Science* **2013**, *342*, 1076. (d) Lin, T. P.; Peters, J. C. *J. Am. Chem. Soc.* **2013**, *135*, 15310. (e) Zhang, G. Q.; Vasudevan, K. V.; Scott, B. L.; Hanson, S. K. *J. Am. Chem. Soc.* **2013**, *135*, 8668.

(3) For a review on Co-catalyzed hydroformylation, see: Hebrard, F.; Kalck, P. *Chem. Rev.* **2009**, *109*, 4272.

(4) For a review on Co-catalyzed hydrovinylation, see: (a) Hilt, G. *Eur. J. Org. Chem.* **2012**, 4441. For recent works, see: (b) Hilt, G.; du Mesnil, F. X.; Lüers, S. *Angew. Chem., Int. Ed.* **2001**, *40*, 387. (c) Grutters, M. M. P.; Müller, C.; Vogt, D. *J. Am. Chem. Soc.* **2006**, *128*, 7414. (d) Hilt, G.; Danz, M.; Treutwein, J. *Org. Lett.* **2009**, *11*, 3322. (e) Arndt, M.; Reinhold, A.; Hilt, G. *J. Org. Chem.* **2010**, *75*, 5203. (f) Sharma, R. K.; RajanBabu, T. V. *J. Am. Chem. Soc.* **2010**, *132*, 3295. (g) Arndt, M.; Dindaroglu, M.; Schmalz, H.-G.; Hilt, G. *Org. Lett.* **2011**, *13*, 6236. (h) Bohn, M. A.; Schmidt, A.; Hilt, G.; Dindaroglu, M.; Schmalz, H.-G. *Angew. Chem., Int. Ed.* **2011**, *50*, 9689. (i) Erver, F.; Hilt, G. *Org. Lett.* **2011**, *13*, 5700. (j) Kersten, L.; Hilt, G. *Adv. Synth. Catal.* **2012**, *354*, 863. (k) Page, J. P.; RajanBabu, T. V. *J. Am. Chem. Soc.* **2012**, *134*, 6556.

(5) For selected reviews on hydroacylation, see: (a) Bosnich, B. *Acc. Chem. Res.* **1998**, *31*, 667. (b) Fu, G. C. In *Modern Rhodium-Catalyzed Reactions*; Evan, P. A., Ed.; Wiley-VCH: New York, 2005; pp 79–91. (c) Jun, C.-H.; Jo, E.-A.; Park, J.-W. *Eur. J. Org. Chem.* **2007**, 1869. (d) Willis, M. C. *Chem. Rev.* **2010**, *110*, 725. (e) Leung, J. C.; Krische, M. J. *Chem. Sci.* **2012**, *3*, 2202.

(6) For selected Rh-catalyzed hydroacylation of alkenes, see: (a) Aloise, A. D.; Layton, M. E.; Shair, M. D. *J. Am. Chem. Soc.* **2000**, *122*, 12610. (b) Jun, C. H.; Lee, D. Y.; Lee, H.; Hong, J. B. *Angew. Chem., Int. Ed.* **2000**, *39*, 3070. (c) Tanaka, M.; Imai, M.; Yamamoto, Y.; Tanaka, K.; Shimowatari, M.; Nagumo, S. J.; Kawahara, N.; Suemune, H. *Org. Lett.* **2003**, *5*, 1365. (d) Willis, M. C.; McNally, S. J.; Beswick, P. J. *Angew. Chem., Int. Ed.* **2004**, *43*, 340. (e) Kundu, K.; McCullagh, J. V.; Morehead, A. T. *J. Am. Chem. Soc.* **2005**, *127*, 16042. (f) Moxham, G. L.; Randell-Sly, H. E.; Brayshaw, S. K.; Woodward, R. L.; Weller, A. S.; Willis, M. C. *Angew. Chem., Int. Ed.* **2006**, *45*, 7618. (g) Stemmler, R. T.; Bolm, C. *Adv. Synth. Catal.* **2007**, *349*, 1185. (h) Tanaka, K.; Shibata, Y.; Suda, T.; Hagiwara, Y.; Hirano, M. *Org. Lett.* **2007**, *9*, 1215. (i) Coulter, M. M.; Dornan, P. K.; Dong, V. M. *J. Am. Chem. Soc.* **2009**, *131*, 6932. (j) Shibata, Y.; Tanaka, K. *J. Am. Chem. Soc.* **2009**, *131*, 12552. (k) Coulter, M. M.; Kou, K. G. M.; Galligan, B.; Dong, V. M. *J. Am. Chem. Soc.* **2010**, *132*, 16330. (l) Phan, D. H. T.; Kou, K. G. M.; Dong, V. M. *J. Am. Chem. Soc.* **2010**, *132*, 16354. (m) Hoffman, T. J.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2011**, *50*, 10670. (n) Murphy, S. K.; Petrone, D. A.; Coulter, M. M.; Dong, V. M. *Org. Lett.* **2011**, *13*, 6216. (o) Vautravers, N. R.; Regent, D. D.; Breit, B. *Commun.* **2011**, *47*, 6635. (p) Chaplin, A. B.; Hooper, J. F.; Weller, A. S.; Willis, M. C. *J. Am. Chem. Soc.* **2012**, *134*, 4885. (q) Murphy, S. K.; Coulter, M. M.; Dong, V. M. *Chem. Sci.* **2012**, *3*, 355. (r) von Delius, M.; Le, C. M.; Dong, V. M. *J. Am. Chem. Soc.* **2012**, *134*, 15022.

(7) For selected Rh-catalyzed hydroacylation of alkynes, see: (a) Kokubo, K.; Matsumasa, K.; Miura, M.; Nomura, M. *J. Org. Chem.* **1997**, *62*, 4564. (b) Tanaka, K.; Fu, G. C. *J. Am. Chem. Soc.* **2001**, *123*, 11492. (c) Jun, C. H.; Lee, H.; Hong, J. B.; Kwon, B. I. *Angew. Chem., Int. Ed.* **2002**, *41*, 2146. (d) Tanaka, K.; Fu, G. C. *J. Am. Chem. Soc.* **2002**, *124*, 10296. (e) Lee, D. Y.; Hong, B. S.; Cho, E. G.; Lee, H.; Jun, C. H. *J. Am. Chem. Soc.* **2003**, *125*, 6372. (f) Tanaka, K.; Fu, G. C. *J. Am. Chem. Soc.* **2003**, *125*, 8078. (g) González-Rodríguez, C.; Pawley, R. J.; Chaplin, A. B.; Thompson, A. L.; Weller, A. S.; Willis, M. C. *Angew. Chem., Int. Ed.* **2011**, *50*, 5134. (h) Lenden, P.; Entwistle, D. A.; Willis, M. C. *Angew. Chem., Int. Ed.* **2011**, *50*, 10657.

(8) For Rh-catalyzed hydroacylation of allenes, see: Osborne, J. D.; Randell-Sly, H. E.; Currie, G. S.; Cowley, A. R.; Willis, M. C. *J. Am. Chem. Soc.* **2008**, *130*, 17232.

(9) For Rh-catalyzed intramolecular hydroacylation of dienes, see: Sato, Y.; Oonishi, Y.; Mori, M. *Angew. Chem., Int. Ed.* **2002**, *41*, 1218.

(10) For Rh-catalyzed hydroacylation of ketones, see: (a) Shen, Z.; Khan, H. A.; Dong, V. M. *J. Am. Chem. Soc.* **2008**, *130*, 2916. (b) Phan, D. H. T.; Kim, B.; Dong, V. M. *J. Am. Chem. Soc.* **2009**, *131*, 15608. (c) Shen, Z.; Dornan, P. K.; Khan, H. A.; Woo, T. K.; Dong, V. M. *J. Am. Chem. Soc.* **2009**, *131*, 1077. (d) Khan, H. A.; Kou, K. G. M.; Dong, V. M. *Chem. Sci.* **2011**, *2*, 407.

(11) For selected Ru-catalyzed hydroacylations, see: (a) Kondo, T.; Akazome, M.; Tsuji, Y.; Watanabe, Y. *J. Org. Chem.* **1990**, *55*, 1286. (b) Kondo, T.; Hiraishi, N.; Morisaki, Y.; Wada, K.; Watanabe, Y.;

Mitsudo, T. *Organometallics* **1998**, *17*, 2131. (c) Shibahara, F.; Bower, J. F.; Krische, M. J. *J. Am. Chem. Soc.* **2008**, *130*, 14120. (d) Omura, S.; Fukuyama, T.; Horiguchi, J.; Murakami, Y.; Ryu, I. *J. Am. Chem. Soc.* **2008**, *130*, 14094. (e) Williams, V. M.; Leung, J. C.; Patman, R. L.; Krische, M. J. *Tetrahedron* **2009**, *65*, 5024. (f) Miura, H.; Wada, K.; Hosokawa, S.; Inoue, M. *Chem.—Eur. J.* **2013**, *19*, 861. (g) Murphy, S. K.; Dong, V. M. *J. Am. Chem. Soc.* **2013**, *135*, 5553.

(12) For Ir-catalyzed hydroacylation, see: Hatanaka, S.; Obora, Y.; Ishii, Y. *Chem.—Eur. J.* **2010**, *16*, 1883.

(13) For selected Ni-catalyzed hydroacylations, see: (a) Tsuda, T.; Kiyoi, T.; Saegusa, T. *J. Org. Chem.* **1990**, *55*, 2554. (b) Taniguchi, H.; Ohmura, T.; Suginome, M. *J. Am. Chem. Soc.* **2009**, *131*, 11298. (c) Hoshimoto, Y.; Ohashi, M.; Ogoshi, S. *J. Am. Chem. Soc.* **2011**, *133*, 4668. (d) Hoshimoto, Y.; Hayashi, Y.; Suzuki, H.; Ohashi, M.; Ogoshi, S. *Angew. Chem., Int. Ed.* **2012**, *51*, 10812.

(14) For selected organocatalyzed hydroacylations, see: (a) Burstein, C.; Glorius, F. *Angew. Chem., Int. Ed.* **2004**, *43*, 6205. (b) Chan, A.; Scheidt, K. A. *J. Am. Chem. Soc.* **2006**, *128*, 4558. (c) Liu, Q.; Perreault, S.; Rovis, T. *J. Am. Chem. Soc.* **2008**, *130*, 14066. (d) DiRocco, D. A.; Oberg, K. M.; Dalton, D. M.; Rovis, T. *J. Am. Chem. Soc.* **2009**, *131*, 10872. (e) Hirano, K.; Biju, A. T.; Piel, I.; Glorius, F. *J. Am. Chem. Soc.* **2009**, *131*, 14190. (f) Biju, A. T.; Wurz, N. E.; Glorius, F. *J. Am. Chem. Soc.* **2010**, *132*, 5970. (g) Bugaut, X.; Liu, F.; Glorius, F. *J. Am. Chem. Soc.* **2011**, *133*, 8130. (h) Liu, F.; Bugaut, X.; Schedler, M.; Fröhlich, R.; Glorius, F. *Angew. Chem., Int. Ed.* **2011**, *50*, 12626. (i) Piel, I.; Steinmetz, M.; Hirano, K.; Fröhlich, R.; Grimme, S.; Glorius, F. *Angew. Chem., Int. Ed.* **2011**, *50*, 4983. (j) Schedler, M.; Wang, D.-S.; Glorius, F. *Angew. Chem., Int. Ed.* **2013**, *52*, 2585.

(15) For Co-catalyzed intermolecular hydroacylations, see: (a) Lenges, C. P.; Brookhart, M. *J. Am. Chem. Soc.* **1997**, *119*, 3165. (b) Lenges, C. P.; White, P. S.; Brookhart, M. *J. Am. Chem. Soc.* **1998**, *120*, 6965. For intramolecular hydroacylation, see: (c) Vinogradov, M. G.; Tuzikov, A. B.; Nikishin, G. I.; Shelimov, B. N.; Kazansky, V. B. *J. Organomet. Chem.* **1988**, *348*, 123.

(16) Fiebig, L.; Kuttner, J.; Hilt, G.; Schwarzer, M. C.; Frenking, G.; Schmalz, H.-G.; Schäfer, M. *J. Org. Chem.* **2013**, *78*, 10485.

(17) For selected other coupling reactions of dienes, see: (a) Shibahara, F.; Bower, J. F.; Krische, M. J. *J. Am. Chem. Soc.* **2008**, *130*, 6338. (b) Smejkal, T.; Han, H.; Breit, B.; Krische, M. J. *J. Am. Chem. Soc.* **2009**, *131*, 10366. (c) Moreau, B.; Wu, J. Y.; Ritter, T. *Org. Lett.* **2009**, *11*, 337. (d) Leung, J. C.; Geary, L. M.; Chen, T.-Y.; Zbieg, J. R.; Krische, M. J. *J. Am. Chem. Soc.* **2012**, *134*, 15700. (e) McInturff, E. L.; Yamaguchi, E.; Krische, M. J. *J. Am. Chem. Soc.* **2012**, *134*, 20628. (f) Zbieg, J. R.; Yamaguchi, E.; McInturff, E. L.; Krische, M. J. *Science* **2012**, *336*, 324. (g) Chen, T.-Y.; Krische, M. J. *Org. Lett.* **2013**, *15*, 2994. (h) Geary, L. M.; Glasspoole, B. W.; Kim, M. M.; Krische, M. J. *J. Am. Chem. Soc.* **2013**, *135*, 3796. (i) Köepfer, A.; Sam, B.; Breit, B.; Krische, M. J. *Chem. Sci.* **2013**, *4*, 1876. (j) Park, B. Y.; Montgomery, T. P.; Garza, V. J.; Krische, M. J. *J. Am. Chem. Soc.* **2013**, *135*, 16320.

(18) For selected cobaltacycle-promoted reactions, see: (a) Chang, H.-T.; Jayanth, T. T.; Cheng, C.-H. *J. Am. Chem. Soc.* **2007**, *129*, 4166. (b) Jeganmohan, M.; Cheng, C.-H. *Chem.—Eur. J.* **2008**, *14*, 10876. (c) Wong, Y.-C.; Parthasarathy, K.; Cheng, C.-H. *J. Am. Chem. Soc.* **2009**, *131*, 18252. (d) Wei, C.-H.; Mannathan, S.; Cheng, C.-H. *J. Am. Chem. Soc.* **2011**, *133*, 6942. (e) Wei, C.-H.; Mannathan, S.; Cheng, C.-H. *Angew. Chem., Int. Ed.* **2012**, *51*, 10592.

(19) Similar metallacycles have been observed in Ni(0)- or Ru(0)-catalyzed cross-coupling of dienes; see refs 17g–i.

(20) For more details, see Supporting Information.

(21) For 2,3-dimethyl-1,3-butadiene, the desired hydroacylation product was obtained in high regioselectivity (>20:1) but with low yield (19%). No coupling product was observed in the hydroacylation of 1,3-cyclohexadiene with benzaldehyde.

(22) Simmons, E. M.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2012**, *51*, 3066.

(23) Blackmond, D. G. *Angew. Chem., Int. Ed.* **2005**, *44*, 4302.

(24) Negishi, E. I.; Huang, Z. H.; Wang, G. W.; Mohan, S.; Wang, C. *Acc. Chem. Res.* **2008**, *41*, 1474.